

## EXPECTED WEED RESPONSE TO PASTURE HERBICIDES

	LATE WINTER TO EARLY SPRING APPLICATIONS							FALL (NOVEMBER TO EARLY DECEMBER) APPLICATIONS					
	2,4-D Ester	2,4-D Amine	ForeFront R&P	Grazon P+D*	Milestone	Redem R&P	Wedmaster	2,4-D Ester	2,4-D Amine	Grazon P+D*	Milestone	Redem R&P	Wedmaster
Bedstraw	3	3	9	9	9	----	3	2	2	9	9	----	2
Broadleaf plantain	8	7	8	8	2	8	9	8	7	8	2	8	9
Buckhorn plantain	7	6	7	8	2	8	8	7	6	8	2	8	8
Bull thistle	8	7	9	9	9	9	8	9	7	9	9	9	9
Buttercups	9	8	9	9	8	9	9	9	8	9	8	9	9
Carolina geranium	5	4	9	9	9	----	7	4	3	9	9	----	6
Common chickweed	2	1	----	8	----	8	5	3	2	9	----	9	8
Curly dock	4	3	9	9	9	----	4	5	3	9	9	----	6
Dandelion	9	8	9	9	9	8	8	9	8	9	9	8	8
Henbit	2	1	----	8	----	8	4	3	2	8	----	8	7
Horseweed	9	8	9	9	9	9	9	9	8	9	9	9	9
Musk thistle	8	7	9	9	9	9	7	9	8	9	9	9	8
Prickly lettuce	8	7	9	9	9	8	8	8	6	9	9	8	8
Red sorrel	3	3	----	8	----	7	5	3	3	8	----	8	6
Sowthistle	8	8	9	9	9	----	9	7	7	9	9	----	8
Wild garlic	8	6	6	5	4	----	6	8	6	5	5	----	6

Key to Response Ratings: 0=No control; 10=100% Control; -- = Data not available

Ratings are based on labeled rates of each herbicide, applied at the optimum time for each weed.

\*For use only in approved TN counties.